

How to read the Atlas

General information

This section provides a brief overview on how to understand the information provided in the species chapters present in this Atlas. In each chapter there is an extended summary of the current state of knowledge about that species, which is aimed to be written in an easily accessible style but at the same time scientifically grounded. Therefore, each chapter has been through a revision by scientific experts and includes a comprehensive list of scientific references. Although the chapters have been written by a number of different authors, they have been harmonized to obtain species information that is as homogeneous as possible throughout the Atlas. The full version of each chapter (expanded and fully peer-reviewed) will be published in the online version of the Atlas at <http://w3id.org/mtv/FISE-Comm/v01/>.

Each chapter starts with a summary and description of the species to continue with paragraphs concerning the species distribution, habitat and ecology, importance and usage and finally threats and diseases. Most chapters deal with a single species, although in a few cases the information is presented at **taxon** level (e.g. circum-Mediterranean firs).

A key contribution of this Atlas is the inclusion of innovative maps and diagrams concerning: 1) Frequency and **Chorology**; 2) Modelled Distribution; 3) Maximum Habitat Suitability; 4) Autoecology, for all those species for which sufficient data exist. High quality images are also included relative to forest habitat, individual trees or more detailed images concerning the bark, leaves, fruits and flowers.

Frequency and Chorology

This map summarises two basic pieces of information concerning the species:

1. The species frequency over a 50 × 50 km grid (blue dots): this shows the percentage of plots inside the grid that contain the species of interest. The sampling points are derived from the same datasets used to model the species distribution (Map 2) and the maximum habitat suitability (Map 3).
2. The species chorology: this is the broad range and qualitative spatial distribution of the tree species derived from one or more bibliographic sources. It is classified as "Native" (green area) when the species is thought to occur naturally and "Introduced" (orange area) when the species has been historically introduced and is nowadays naturalised. In cases where it is not possible to distinguish between the natural and introduced range, "Actual" range is shown.

For more details on the datasets and methodology used, as well as a list of the bibliographic sources used to construct the chorology, see the chapter "modelling, data and information on forest tree species" on page 40.

Technical terms

Technical words are presented in **this font** and are listed in the glossary on page 190 at the end of the Atlas.

Fagus sylvatica

Fagus sylvatica in Europe: distribution, habitat, usage and threats


T. Houston Durrant, D. de Rigo, G. Caudullo

Fagus sylvatica L., or European beech, is one of the most important and widespread broadleaved trees in Europe. It is a large deciduous tree that can maintain its high growth rate until late maturity. Its natural range extends from southern Scandinavia to Sicily, from Spain in the west to northwest Turkey in the east. Though not demanding of soil type, beech requires a humid atmosphere with precipitation well distributed throughout the year and a well-drained soil. It tolerates rigorous winter cold, but is sensitive to spring frost. Owing to the capacity of its root system for assisting in the circulation of air throughout the soil, and the amount of potash in its leaves, Beech trees conserve the productive capacity of the soil better than many other species. Its wood is strong and wears well making it ideal for a wide range of uses, from furniture to musical instruments, as well as for pulp and firewood.

The European beech (*Fagus sylvatica* L.) is a large deciduous tree that commonly reaches 30-40m and is capable of attaining heights up to 50m in some locations¹. In contrast to many other tree species, it is able to maintain a high rate of growth until a relatively mature age. The tree is usually single-stemmed with silver-grey bark. The leaves are typically 10 × 7 cm, dark and shiny green. They have an oval to elliptic shape, with wavy margins and the end of the parallel veins on each side^{2,3}. Beech is monoecious: the male and female flowers are borne on the same tree. It has a typical life span of around 150-300 years, and reproduces very late (40-50 years old). Fruiting normally occurs every 5 to 8 years. Its seed production is characterised by irregular mast years (when a very heavy crop is produced), usually following hot summers of the previous year. The bitter edible nuts are sharply tri-angled and are borne singly or in pairs in soft-spined husks. The beech nuts are an important source of food for several animals; and birds including squirrels, woodpigeons, woodpeckers and jays; they also play a major part in seed dispersal by hiding the seeds and falling to retrieve all of them⁴.


Distribution

Beech is widespread across Europe: it can be found from Sicily in the south to Bergen in southern Norway^{5,6}. An analysis of pollen records indicate that the species has spread across Europe from small scattered populations left after the last glaciation, and is currently probably at its maximum post-glacial spread⁷. It needs a growing season of at least 140 days, and for this reason cannot survive too far north in Scandinavia⁸. Longitudinally its range is from the Cantabrian Mountains in the west to the Carpathians and Balkan Mountains in the east, although there are some areas in Europe where it is not found as a native tree, such as the Po valley and the Hungarian plain. As the climate becomes more continental in the eastern parts of Europe it is replaced by oriental beech (*Fagus orientalis*).




Map 1: Plot distribution and simplified chorology map for *Fagus sylvatica*. Frequency of *Fagus sylvatica* occurrences within the field observations as reported by the National Forest Inventories. The chorology of the native spatial range for *F. sylvatica* is derived after Meusel and Jäger, and EURFORGEN^{19,20}.

At the southern part of its range (Spain, Sicily) it is only normally present at altitudes of more than 1000m, and can even be found at elevations of up to 2000m^{1,8}. High summer temperatures, drought and moisture availability are limiting factors for the distribution of beech in Europe, but continentality is also associated with limiting its spread in north-western regions⁴. Climate change may have impacts on its future distribution, particularly at the extremes of its range where it is likely to become less competitive in the south and east (primarily because of drought), but could expand its range into Scandinavia and the Baltic⁹.



Map 2: High resolution distribution map estimating the relative probability of presence.



Large beech in a mountain pasture in Piani di Praglia (Genova, North Italy). Copyright Ettore Balestracci, www.fidocom.it/01/

Habitat and Ecology

Beech is a hardy species. It tolerates very shady situations (it is the most shade-tolerant broadleaved tree in its range¹⁰), so that natural regeneration is possible in silvicultural systems with continuous crown coverage as the seedlings are able to survive and grow below the canopy of established trees. The predominance of beech means a reduction of light level in the understorey vegetation level and in that case beech seeds survive better than those of other tree species. It is not particularly soil-sensitive¹¹ and grows on a wide variety of soils with a pH range from 3.5 to 8.5, although it cannot tolerate the most acidic conditions. Beech shows a moderate soil-acidifying ability¹². It prefers moderately fertile ground, calcified or lightly acidic and is also sensitive to late frosts¹³, therefore it is found more often on the side of a hill than at the bottom of a clayey basin. It grows well on soft soils in which the root system can easily penetrate and its optimal growth is in humid soils situated on calcareous or volcanic parent rocks. On the contrary, it does not thrive on sites that are regularly flooded or which have stagnant water, since it needs good drainage and will not tolerate waterlogged or compacted soils^{1,14}. Beech furthers soil conservation due to its production of a large quantity of litter (around 900g/m² per year). The root system tends to be shallow, making it susceptible to drought when compared to coniferous stands¹⁵. However, there appears to be some genetic variability across different climatic zones, since trees in southern Europe are able to cope better with drought than those in the north¹.

Importance and Usage

Beech is an important European forestry tree. Fine grained and knot-free, the wood is hard and has a pale cream colour and good workability¹⁶. With around 250 known usages, it is one of the most diversely used tree species in Europe. Its wear-resistance, strength, and excellent bending capabilities make it ideal for boatbuilding, flooring, stairs, furniture, musical instruments (piano pinblocks), plywood, panels, veneering and cooking utensils such as bowls, platters and wooden spoons. It is also used for pulp and can be coppiced for fire wood and charcoal due to its relatively high energetic potential^{1,9,16}.

Threats and Diseases

The root system architecture of beech may vary depending on local soil conditions¹⁷. While generally showing a noticeable resistance to rockfall and wind-throw^{12,18}, under unfavourable local conditions a relatively shallow root system may make the tree vulnerable to wind-throw⁴. The thin bark provides little



Shiny dark green leaves with red galls caused by the fly Mikotoz fagi (Diptera Cecidomyiidae). Copyright AR60002, commons.wikimedia.org CC0

Modelled spatial distribution of the species

This map represents the **relative probability of presence** of the species derived from a harmonised dataset of forest **field observations** made from a number of different surveys and available within the Forest Information System for Europe (FISE). The maps are presented at a high-spatial resolution of 1 km. Dark green colour means the species is very likely to be found at that location, while the pale brown colour signifies a low probability of presence. For some regions there were

not enough data available to make any predictions about the probability of presence; these are coloured pale grey. The map has been modelled with an innovative methodology designed to take into account the different local densities of the underlying data sets.

For more details on the data set and modelling techniques used, see the Atlas chapter "modelling, data and information on forest tree species" on page 40.

Key Fact

In some chapters there is a supplementary box focusing on some particular aspects of the species or taxon considered (e.g. notes on the taxonomy of the species or information about a related species).


Fagus sylvatica

Fagus orientalis

Fagus orientalis, or oriental beech, is closely related to *Fagus sylvatica*. Some authorities consider them to be sub-species; others consider them to be two separate species¹. In appearance they are generally very similar. The leaves are slightly longer, darker and less glossy than those of European beech, and tend to have more vein-pairs (9-14 as opposed to 5-9)². Oriental beech can be found in the Balkans, Anatolia, the Caucasus, northern Iran and Crimea³. Its range overlaps with that of the European beech and there is frequently hybridisation between the two⁴. Where both species are present, oriental beech tends to favour the valleys while European beech is found further up the slopes; this is because the European beech is more susceptible to late frosts⁵.




Foliage and fruits of oriental beech (*Fagus orientalis*).
Copyright: DidiKub, commons.wikimedia.org, CC BY



protection from fire, and can also be damaged through stripping and gnawing by squirrels and other mammals. The presence of deer is a limiting factor because they eat young stands. Spring frosts often damage young trees or flowers appearing at the same time as leaves. Young beech trees are susceptible to woolly aphid; mature trees can suffer internal rot by the fungus *Ganoderma applanatum*. Old trees (100-1200 years) may suffer 'red heart' which reduces stability and timber value⁶. Beech is among the susceptible hosts to *Phytophthora ramorum* and



Mature beech forest with autumn colour foliage in Delamere forest, Cheshire, UK.
Copyright: Freepress, www.forest.gov.uk, © Crown Copyright



Field data in Europe (including absences) ● Observed presences in Europe ● Autoecology diagrams based on harmonised field observations from forest plots.

large regions across Europe have climatic suitability to this pest, which may become a more serious problem in the future⁷. The large pine weevil (*Hyllobius abietis*) is harmful to beech and markedly coexists with part of its natural niche. Herbivory by short-snouted weevils (*Strophosoma melanogrammum* Forst. and *Otiorhynchus scaber*) is another threat to beech^{8,9}.

References

- [1] J. R. Pochham, P. A. Thomas, M. D. Atkinson, T. Diegel, *Journal of Ecology* **100**, 1557 (2012).
- [2] A. F. Mitchell, P. Dahlstrom, E. Suresen, C. Chanters, *Field guide to the trees of Britain and northern Europe* (Collins, 1974).
- [3] O. Johnson, D. More, *Collins tree guide* (Collins, 2005).
- [4] J. Fang, M. J. Lachowicz, *Journal of Biogeography* **33**, 1804 (2006).
- [5] R. Baker, et al., *EFSA Journal* **9**, 2186+ (2011).
- [6] E. Hulthén, M. Fries, *Atlas of North European vascular plants (North of the Tropic of Cancer)*, Vols. I-III. (Koeltz scientific books, 1966).
- [7] D. Magri, *Journal of Biogeography* **35**, 450 (2008).
- [8] T. Horgan, et al., *A guide to forest tree species selection and silviculture in Ireland*. (National Council for Forest Research and Development (COFORD), 2003).
- [9] K. Kramer, et al., *Forest Ecology and Management* **259**, 2213 (2010). The ecology and silviculture of beech: from gene to landscape.
- [10] A. Pratiak, et al., *The CABI encyclopedia of forest trees* (CABI, Oxfordshire, UK, 2013).
- [11] L. Walther, E. Graf Pannatier, E. S. Meier, *Forest Ecology and Management* **297**, 94 (2013).
- [12] L. Augusto, J. Ranger, D. Binkley, A. Rothe, *Annals of Forest Science* **59**, 233 (2002).
- [13] L. Paule, *Forest Genetics* **2**, 161 (1995).
- [14] A. Geller, et al., *Trees - Structure and Function* **21**, 1 (2007).
- [15] A. Granier, et al., *Agricultural and Forest Meteorology* **143**, 123 (2007).
- [16] M. Goldstein, G. Simonetti, M. Watschinger, *Atlas of Europe's Flora* (Koenigstein, 1992).
- [17] J. E. Norris, A. Di Iorio, A. Stokke, B. C. Nicoli, A. Achim, *Slope Stability and Erosion Control: Ecotechnological Solutions*, J. E. Norris, et al., eds. (Springer Netherlands, 2008), pp. 187-210.
- [18] L. K. A. Doreen, F. Berger, C. de Hér, E. Mermin, P. Taroff, *Forest Ecology and Management* **215**, 183 (2005).
- [19] J. I. Barredo, et al., *EPPO Bulletin* **45**, 275 (2015).
- [20] CABI, *Hyllobius abietis* (Large pine weevil) (2013). *Invasive Species Compendium*, 134, 111 (2000).
- [21] M. Löf, G. Isacson, D. Rydberg, T. N. Wälinder, *Forest Ecology and Management* **190**, 281 (2004).
- [22] G. Karamic, Z. Kaya, *EUFORGEN Technical guidelines for genetic conservation and use for Oriental beech (Fagus orientalis)* (Biodiversity International, Rome, Italy, 2009).
- [23] A. Oprea, C. Sibiu, L. Gola, *Contributi Botanice* **46**, 17 (2011).
- [24] M. Štajčič Nikolić, J. Milovanović, M. Nosić, R. Knežević, D. Štanić, *Genetika* **45**, 369 (2013).
- [25] T. Denk, G. Grimm, K. Stögerer, M. Langer, V. Hemleben, *Plant Systematics and Evolution* **232**, 215 (2002).
- [26] H. Meusel, E. J. Jäger, *Plant Systematics and Evolution* **162**, 315 (1989).
- [27] EUFORGEN, *Distribution map of beech (Fagus sylvatica)* (2008). www.euforgen.org
- [28] J. Jalas, J. Suominen, *Atlas Florae Europaeae: distribution of vascular plants in Europe Vol. 3: Salicaceae to Balsanifloraceae* (Committee for Mapping the Flora of Europe and Societas Biologica Fennica Vanamo, Helsinki, 1976).
- [29] K. Brownicki, J. Zielinski, *Chronology of trees and shrubs in south-west Asia and adjacent regions*, vol. 1 (Polish Scientific Publishers, Warszawa, 1976).

This is an extended summary of the chapter. The full version of this chapter (revised and peer-reviewed) will be published online at <https://w3id.org/mtv/FISE-Comm/v01/012690>. The purpose of this summary is to provide an accessible dissemination of the related main topics. This QR code points to the full online version, where the most updated content may be freely accessed. Please, cite as: T. Houston Durrant, D. de Rigo, G. Caudullo, 2016. *Fagus sylvatica* and other beeches in Europe: distribution, habitat, usage and threats. In: European Atlas of Forest Tree Species. e012690+. Publ. Off. EU.



Tree species | European Atlas of Forest Tree Species 95

Modelled maximum habitat suitability

This map represents the Maximum Habitat Suitability of the species, namely the areas where the species could potentially occur if climatic conditions and ecological conditions are met. It is modelled based on a harmonised and very dense dataset of forest plots available for most of Europe (see Map 1, Frequency) in combination with high-resolution bioclimatic parameters (e.g. temperature and precipitation), solar irradiation and elevation range. The maps are presented at a high-spatial resolution of 1 km.

Dark blue areas represent areas that are highly suitable for the species to survive (denoted in the legend as high survivability areas). In these areas, the local bioclimatic conditions are very similar to those of at least some of the field observations where the species occurs. Conversely, orange areas highlight low survivability conditions. This refers to areas with a bioclimatic pattern very dissimilar from all the observed patterns where the species is found.

In practice, a given species may not be found in all the areas marked "high survivability" for other reasons (e.g. competition from or preferential planting of other species). However, in those areas marked "negligible survivability" the species is unlikely to grow, even if deliberately planted there.

The map is modelled with an innovative methodology taking into account the different spatial distributions of the underlying datasets as well as a number of bioclimatic and geographic factors. Detailed information on the data and techniques used can be found in the chapter "modelling, data and information on forest tree species" on page 40.

Images

Images have been carefully selected to help identification/understanding of species. Captions also identify the individuals who have provided the image and the relative copyrights. A complete list of contributors is at the beginning of the atlas.

References

Each chapter has been fully referenced with the most up-to-date scientific literature, which has been revised by three scientific experts. All the references are sequentially included through the text and listed at the end of the chapter. A revised and extended set of references will be available in the full online version of each chapter.

QR chapter code and citation information

This code points to the full online version of the species chapter, where the most up-to-date content can be found.

The correct way to cite this extended summary is also shown here (in the online version, readers will find the correct way to cite the full chapter).

In the online version it will additionally be possible to navigate through the expanded, fully peer-reviewed version of the chapter having the possibility to download maps, diagrams and text.

The online version will be hosted within the newly established Forest Information System for Europe (FISE). The Online European Atlas of Forest Tree Species will be part of the FISE Communications (FISE-Comm):

<http://w3id.org/mtv/FISE-Comm/v01/>

Autoecology Diagrams

In most chapters, autoecology diagrams (also known as climate-space diagrams) have been derived for the described species, based on the datasets of field observations as harmonised within the Forest Information System for Europe (FISE). These observations are the same as those used to estimate the coarse-resolution forest plot distribution presented in Map 1. The local bioclimatic conditions where a given species is observed are obtained by means of a number of high-resolution bioclimatic and geographic variables. The number of

possible combinations of variables is very large and for this Atlas we have focussed on three: 1. Annual average temperature vs. Annual precipitation; 2. Potential spring-summer solar irradiation vs. Average temperature of the coldest month; 3. Seasonal variation of monthly precipitation vs. Sum of precipitation of the driest month. In the online version of the Atlas other combinations of variables may also be found.

The overall climate space occupied by each of the field observations on every species is represented by a grey

spot (one for every plot), while those plots containing the species of interest are coloured blue, thus illustrating the specific climate niche of that species, and showing how a given species might be constrained by one or more climatic conditions. Grey patches on this page may be coloured blue in others where different species are adapted to different conditions.

For more details on the data and modelling aspects, see the Atlas chapter "modelling, data and information on forest tree species" on page 40.