

## Salix caprea in Europe: distribution, habitat, usage and threats

C. M. Enescu, T. Houston Durrant, D. de Rigo, G. Caudullo

*Salix caprea* L., commonly known as goat willow, is a pioneer and a fast-growing plant, which has a wide distributional area across Europe and Asia in the boreal and temperate zones. The scientific name *caprea* means goat, which probably derives from the fact that its leaves were used as goat fodder. Due to its wide distribution range and its high ecological amplitude, goat willow represents a very valuable multi-purpose species, used principally for biomass plantation and for gardening and hedges.


The goat willow (*Salix caprea* L.) is a deciduous small to medium-size tree or shrub with heights up to 10m<sup>1, 2</sup>, reaching only exceptionally 15m<sup>3</sup>. It has a far-reaching and especially well-developed fibrous root system<sup>1</sup>. The trunks are small, occasionally reaching a diameter of 40cm<sup>4</sup>. The bark is smooth at the beginning, forming a thin **rhytidome**, irregularly cracked<sup>2</sup>. The twigs are greenish, thick and with grey hairs. The leaves are alternate, broadly elliptic and up to 5-12cm long. The upper side of the leaf is green and glossy, while the underside is covered by densely and softly white-downy hairs<sup>1, 5</sup>. The goat willow is a **dioecious** species and catkins appear in early spring (usually in March or April, depending on the site conditions). Male catkins have spreading yellow stamens, while female catkins are greenish and insect-pollinated. The catkins are erect, with approximately 100-200 flowers in each female catkin and 200-300 flowers in each male catkin, respectively. Catkins are produced abundantly and they appear before the leaves<sup>3</sup>. This willow naturally hybridises with a number of other *Salix* species, producing fertile descendants with intermediate characteristics. Hence, this makes it difficult to determinate taxonomical limits and to assign individuals to a particular species within the *Salix* genus<sup>6, 7</sup>.

### Distribution

This species is native to cool temperate and boreal regions of Europe and Asia, occurring in a large range of habitats across Europe and Asia<sup>8</sup>. Its wide distribution spreads from Spain to China, from Turkey and northern Iran to **Fennoscandia** and Siberia reaching up to 70° north latitude<sup>1, 9-11</sup>. It has been also introduced in eastern North America and is now naturalised<sup>12</sup>.

### Habitat and Ecology

The goat willow is a pioneer<sup>13</sup> and a fast-growing plant<sup>14</sup>, tolerating a wide variety of **edaphic** and climatic conditions<sup>1</sup>. It shows a high adaptability to different habitat conditions<sup>15</sup>, such as dry or wet sites<sup>16</sup>, usually growing in **mesic** to moist stands<sup>4</sup>. It prefers the calcareous sites, but it grows on almost all soil



Map 1: Plot distribution and simplified chorology map for *Salix caprea*. Frequency of *Salix caprea* occurrences within the field observations as reported by the National Forest Inventories. The chorology of the native spatial range for *S. caprea* is derived after several sources<sup>9, 52-56</sup>.

types<sup>17</sup>. Compared with other willow species, the goat willow is more sensitive to continuous flooding<sup>18</sup>, avoiding the saturated soils of wetlands<sup>19</sup>. Regarding its demand for light, it is a very shade intolerant species, occurring in open areas with full sun<sup>1</sup>, but it is one of the few willow species able to grow in forest understories<sup>8</sup>. Unlike almost all other willows, the goat willow can only occasionally be propagated vegetatively by cuttings<sup>8, 20</sup>.

In boreal forests, it is found in the dominant Norway spruce (*Picea abies*), Scots pine (*Pinus sylvestris*) and birch (*Betula* spp.) forests, in admixture with other deciduous species such as aspen (*Populus tremula*) and rowan (*Sorbus aucuparia*)<sup>21, 22</sup>, favoured by disturbances which create open areas<sup>23</sup>. In temperate forests dominated by oaks (*Quercus* spp.), European beech (*Fagus*

*sylvatica*) and Norway spruce, it is present with other understorey species such as hazel (*Corylus avellana*), elder (*Sambucus nigra*) and bramble shrubs of the genus *Rubus*, occurring principally in forest clearings and open areas<sup>24-26</sup>.




Goat Willow in Hesse, Germany. (Copyright Willow, commons.wikimedia.org. CC-BY)

### Importance and Usage

This species has a wide variety of uses. The scientific name *caprea* actually means goat, which probably derives from the fact that its leaves are used as fodder<sup>27</sup>. In northern Europe, it is a common species in the agricultural landscape, used as a windbreak and hedge<sup>28</sup>, and its foliage is used as fodder for cattle and goats<sup>29, 30</sup>. In traditional medicine, goat willow extracts are used as a painkiller, astringent, antiseptic, eye tonic<sup>28</sup>, or even to treat malaria, gout, neuralgia and intestinal diseases<sup>31</sup>. As is also the case with other willows, a good charcoal can be obtained from its burnt wood, which in the past was used for gunpowder and for drawing pencils<sup>32, 33</sup>. The goat willow is particularly valuable for its high biomass production in short rotation plantations and its role in landscape restoration<sup>34</sup>, as it is able to tolerate even polluted land<sup>35</sup>. Worldwide, this willow is also used for its ability to extract heavy metals, such as Cd and Zn, (i.e. phyto-extraction strategy) from polluted sites<sup>34-36</sup>. During spring holidays, in several places across Europe, its flowering branches are in high demand<sup>37</sup>. It is also appreciated as a **melliferous** plant species<sup>38, 39</sup>, supplying a honey production of 150-200 kg/ha<sup>40</sup>. The goat willow distribution range overlaps with many areas in Europe with high erosion rates, including European boreal areas and moist slopes with high drainage-area within the European mountain systems<sup>41</sup>. In these critically erosion-prone areas, it contributes to key ecosystem services such as watershed protection and soil stabilisation<sup>42</sup>. As for other willows and poplars, it is also useful for ecosystem restoration and **phytoremediation**<sup>43, 44</sup>.

Finally, the goat willow plays an important role in maintaining species diversity, by being the host of several lichen species<sup>4, 37</sup>.



Map 2: High resolution distribution map estimating the relative probability of presence.



Male catkins turn yellow when the pollen is ready for release. (Copyright AnRo002, commons.wikimedia.org. CC0)

This willow is among the first flowering plants in the spring, the catkins providing a high quantity of pollen and nectar as a food source, to bees, insects or birds, such as Eurasian blue tits (*Cyanistes caeruleus*)<sup>3</sup>.



Female catkins are insect pollinated. (Copyright AnRo0002, commons.wikimedia.org, CC0)




## Threats and Diseases

The goat willow is a short-living plant and has no severe threatening diseases in its natural habitats. Pests are reported to result in economic loss only in biomass coppice plantations. The fungus *Rhytisma salicinum*, the leaf-galler sawfly *Pontania pedunculii* and the leaf-folder sawfly of genus *Phyllocolpa* may cause damage to individuals, by affecting the leaves<sup>45</sup>. The rust fungus *Melampsora capaeorum* may infect the goat willow and its hybrids, raising dusty orange spots or pustules on the leaves, resulting in lower photosynthetic performance and eventually defoliating the trees. Breeding programmes have been promoted for selecting rust resistant willows in biomass plantations and gardening plants<sup>46, 47</sup>. The goat willow is susceptible to attacks



The leaves are broader than those of many other willow species. (Copyright AnRo0002, commons.wikimedia.org, CC0)

Field data in Europe (including absences) ● Observed presences in Europe ●



Autoecology diagrams based on harmonised field observations from forest plots.

## References

- [1] F. Clinovschi, *Dendrologie* (Editura Universitatii Suceava, 2005).
- [2] I. D. Trava, G. F. Borlea, W. Hollerbach, *Journal of Horticulture, Forestry and Biotechnology* **18**, 209 (2014).
- [3] Q. O. N. Kay, *Bird Study* **32**, 40 (1985).
- [4] M. Kuusinen, *Mitt. Eidgenöss. Forsch. anst. Wald. Schnee. Landsch.* **70**, 125 (1995).
- [5] B. Eversham, *A key to lowland willows, sallows and osiers* (2006). V. 1.6.
- [6] M. Aránzazu Prada, D. Arizpe, *Riparian Tree and Shrub Propagation Handbook: An Aid to Riverine Restoration in the Mediterranean Region* (Generalitat Valenciana, Valencia, 2008).
- [7] A. E. Palme, V. Semerikov, M. Lascoux, *Heredity* **91**, 465 (2003).
- [8] H. Neuner, R. Beiderbeck, *Silvae Genetica* **42**, 308 (1993).
- [9] H. Meusel, E. Jäger, S. Rauschert, E. Weinert, *Vergleichende Chorologie der Zentraleuropäischen Flora* (Gustav Fischer Verlag Jena, 1978).
- [10] E. Hultén, M. Fries, *Atlas of North European vascular plants (North of the Tropic of Cancer)*, Vols. I-III. (Koeltz scientific books, 1986).
- [11] J. Jalas, J. Suominen, *Atlas Florae Europaeae: distribution of vascular plants in Europe Vol. 3 Salicaceae to Balanophoraceae* (Committee for Mapping the Flora of Europe and Societas Biologica Fennica Vanario, Helsinki, 1976).
- [12] G. W. Argus, *Flora of North America North of Mexico*, Flora of North America Editorial Committee, ed. (New York and Oxford, 2010), vol. 7.
- [13] A. Bobiec, *Polish Journal of Ecology* **55**, 441 (2007).
- [14] J. B. Faliński, *Journal of Vegetation Science* **9**, 57 (1998).
- [15] J. Dušek, J. Květ, *Biologia* **61**, 441 (2006).
- [16] T. F. Marshall, *Section 5: Salicaceae (willows & poplars)* (Prestwood Nature, 2014).
- [17] S. Lowe, N. Repper, L. Miles, S. G. Wallace, *Notes on tree planting and the use of native species in North East England* (Wildlife Trusts, Northumberland, UK, 2012).
- [18] R. J. Talbot, J. R. Etherington, J. A. Bryant, *New Phytologist* **105**, 563 (1987).
- [19] J. G. Isebrands, J. Richardson, *Poplars and willows: trees for society and the environment* (CABI; FAO, 2014).
- [20] M. Liesebach, G. Naujoks, *Plant Cell, Tissue and Organ Culture* **79**, 239 (2004).
- [21] E. Lähde, T. Eskelinen, A. Väänänen, *Forestry* **75**, 395 (2002).
- [22] T. Kuuluvainen, *Silva Fennica* **36** (2002).
- [23] O. Engelmarm, *Boreal forest disturbances* (Elsevier, 1999), chap. 6, pp. 161–186.
- [24] M. Budzaková, D. Galvánek, P. Littera, J. Šibík, *Acta Societatis Botanicorum Poloniae* **82**, 13 (2013).
- [25] R. Van Couwenbergh, C. Collet, E. Lacombe, J.-C. Pierrat, J.-C. Gégout, *Forest Ecology and Management* **260**, 146 (2010).
- [26] T. Degen, F. Devillez, A.-L. Jacquemart, *Annals of Forest Science* **62**, 429 (2005).
- [27] J. E. Smith, *The English Flora Vol. 4* (Longman, Rees, Orme, Brown and Green, London, 1830), second edn.
- [28] A. Ahmed, W. A. Shah, S. Akbar, M. Younis, D. Kumar, *International Journal of Research in Phytochemistry & Pharmacology* **1**, 17 (2011).
- [29] A. Popp, K.-M. Scheibe, *Agriculture* **3**, 147 (2013).
- [30] J. Smith, *The History of Temperate Agroforestry* (The Organic Research Centre, Elm Farm, Newbury, Berkshire, UK, 2010).
- [31] A. Ahmed, et al., *Journal of Pharmacy Research* **4**, 1067 (2011).
- [32] F. P. Porcher, *Resources of the southern fields and forests, medical, economical, and agricultural* (Walker, Evans & Cogswell, Charleston, South Carolina, U.S., 1869).
- [33] W. Withering, *A botanical arrangement of all the vegetables naturally growing in Great Britain*, vol. 2 (Cadel and Elmsley, London, 1776).
- [34] M. Vaculik, et al., *Environmental Pollution* **163**, 117 (2012).
- [35] C. Varga, M. Marian, L. Mihaly-Cozmuta, A. Mihaly-Cozmuta, L. Mihalescu, *Analele Universitatii din Oradea, Fascicula Biologie* **16**, 141 (2009).
- [36] M. Kuffner, et al., *Journal of Applied Microbiology* **108**, 1471 (2010).
- [37] W.-D. Gu, M. Kuusinen, T. Kontinen, I. Hanski, *Ecography* **24**, 139 (2001).
- [38] N. Ion, *International Scientific Symposium* (University of Agronomical Sciences and Veterinary Medicine of Iasi, Romania, 2008).
- [39] A. C. Băloiu, I. Csoș, S. C. Martin, A. Bogluț, *Lucrări Științifice. Management Agricol* **15**, 241 (2013).
- [40] D. C. Popovici, E. Crăniceanu, D. S. Bistrea, O. Colibar, *Lucrări Științifice Medicină Veterinară* **40**, 46 (2007).
- [41] C. Bosco, D. de Rigo, O. Dewitte, J. Poesen, P. Panagos, *Natural Hazards and Earth System Science* **15**, 225 (2015).
- [42] J. E. Norris, A. Di Iorio, A. Stokes, B. C. Nicoll, A. Achim, *Slope Stability and Erosion Control: Ecotechnological Solutions*, J. E. Norris, et al., eds. (Springer Netherlands, 2008), pp. 167–210.
- [43] Y. Kuzovkina, M. Quigley, *Water, Air, and Soil Pollution* **162**, 183 (2005).
- [44] J. Ball, J. Carle, A. Del Lungo, *Unasylva* **56**, 3 (2005).
- [45] J. Hjäältén, L. Ericson, H. Roininen, *Ecoscience* **7**, 51 (2000).
- [46] B. Buhner, et al., *Fungal Biology* **118**, 910 (2014).
- [47] M. H. Pei, T. Hunter, D. J. Royle, *New Phytologist* **141**, 155 (1999).
- [48] D. de Rigo, et al., *Scientific Topics Focus* **2**, m110a15+ (2016).
- [49] V. de Tillesse, L. Nef, J. Charles, A. Hopkin, S. Augustin, *Damaging poplar insects - Internationally important species* (International Poplar Commission, FAO, Rome, 2007).
- [50] M. J. Lechowicz, Y. Mauffette, *Revue d'Entomologie du Québec* **31**, 43 (1986).
- [51] E. M. Andreeva, *Russian Journal of Ecology* **33**, 342 (2002).
- [52] Botanical Society of Britain & Ireland, *BSBI big database* (2015). <http://bsbidb.org.uk>
- [53] B. Jonsell, ed., *Flora Nordica, Vol. 1: Lycopodiaceae - polygonaceae* (The Royal Swedish Academy of Sciences, Stockholm, 2000).
- [54] A. K. Skvortsov, *Willows of Russia and Adjacent Countries: Taxonomical and Geographical Revision* (English translation of 1968 Russian edition) (University of Joensuu, Finland, 1999).
- [55] Anthos, *Information System of the plants of Spain* (Real Jardín Botánico, CSIC - Fundación Biodiversidad, 2015). <http://www.anthos.es>.
- [56] Tela Botanica, eFlora (2015). <http://www.tela-botanica.org>

This is an extended summary of the chapter. The full version of this chapter (revised and peer-reviewed) will be published online at <https://w3id.org/mtv/FISE-Comm/v01/e01322d>. The purpose of this summary is to provide an accessible dissemination of the related main topics.

This QR code points to the full online version, where the most updated content may be freely accessed. Please, cite as:  
Enescu, C. M., Houston Durrant, T., de Rigo, D., Caudullo, G., 2016. *Salix caprea* in Europe: distribution, habitat, usage and threats. In: San-Miguel-Ayanz, J., de Rigo, D., Caudullo, G., Houston Durrant, T., Mauri, A. (Eds.), *European Atlas of Forest Tree Species*. Publ. Off. EU, Luxembourg, pp. e01322d+

